

Most commonly anorexia & bulimia nervosa (prevalence ~1-2% each, >90% are F) which peak in 2<sup>nd</sup>-3<sup>rd</sup> decades and are characterised by disturbances in thinking & behaviour re food, eating & body weight or shape. Other eating disorders include morbid obesity and pica.

*Feature of AN & BN include:*

- Distorted body image
- Self-induced vomiting, misuse of laxatives, diuretics or appetite suppressants
- Excessive exercise
- In anorexia nervosa:
  - Morbid fear of weight gain/fatness
  - Restricted dietary intake
  - Amenorrhoea or failure to reach menarche in women, loss of sexual interest in men
  - BMI ≤ 17.5 for adults with wt ≤ 85% normal.
- In bulimia nervosa:
  - Preoccupation with food, weight and shape
  - Cycles of binge eating episodes, followed by purging, excessive dieting or exercise
  - Weight/BMI tends to be normal-slightly overweight.

Comorbid psychiatric illnesses are common (up to 80%) including: depression, anxiety disorders, OCD, substance abuse, DSH & suicidal ideation

*Complications:* Malnutrition, electrolyte imbalance esp hypokalaemia, sepsis, RF, bone marrow suppression, acute oesophageal tears, amenorrhoea, constipation, suicide attempts.

## Management

### Safety

### Assessment

### Confirmation of provisional diagnosis

### Consultation

### Immediate treatment

### Transfer of care

## Safety

Observation is important as patients may try to purge themselves in ED. Also a suicide risk.

## Assessment

Often poor insight, denial, reluctance to give history & non-compliance with management.

### History:

Attitudes to food. Eating, weight gain pattern. Socio-cultural background. Family dynamics.

- *The SCOFF Questionnaire (>1 positive response → possible disorder)*
  - Do you ever make yourself sick because you feel uncomfortably full?
  - Do you worry you have lost control over how much you eat?
  - Have you recently lost more than 6kg in a three month period?
  - Do you believe yourself to be fat when others say you are too thin?
  - Would you say that food dominates your life?

### Exam - Physical & Mental State

Physical exam may reveal emaciation, lanugo hair. Also pitted teeth, parotid swelling & Russell's sign (calluses on knuckles) from repeated induced vomiting. Evidence of Cx: dehydration, sepsis, arrhythmias (bradycardia), hypothermia, anaemia, osteoporosis, acute gastric dilation (binging).

In MSE screen for evidence of depression, anxiety, obsessionality, substance abuse, or other psychiatric condition, DSH; ask about suicide ideation and assess insight/motivation.

### Confirmation of provisional diagnosis

Corroboration: Important as patients often deny there is a problem. Family, friends, GP, notes.  
Investigations: Weight/BMI, U/A, ECG, UEC, CMP, BSL, FBC, TFT, LFT, ABG, hormone levels.

### Consultation

Mental health± Adolescent Health teams.

### Immediate treatment

Rehydration, electrolyte replacement, nutritional support. Rx for complications.  
D/W MH team re combination RX of antidepressants & CBT.

### Transfer of care

Consider admission if on medical grounds if:

- $T < 35.5^{\circ}\text{C}$
- $\text{BP} < 90/60\text{mmHg}$  in adults or  $< 80/40\text{mmHg}$  in adolescents or postural drop  $\geq 20\text{mmHg}$
- Tachycardia / bradycardia or other ECG abnormality
- $\text{BMI} < 15\text{kg/m}^2$  or rapid weight loss ( $\geq 1\text{kg/week}$  over  $> 4\text{w}$ )
- Significant electrolyte disturbance (e.g.  $\text{K}^+ < 3.0\text{mmol/L}$ )
- Dehydration or urinary ketones
- Unable to eat
- Refeeding syndrome

Consider admission if on mental health grounds if:

- Suicidal ideation or significant eating disordered symptoms
- Treatment resistance or sabotage
- Treatment of comorbid mental illness

Discharge may be possible after consultation with MH team and referral to appropriate community-based eating disorders services, a mental health professional, GP and dietician.